



ELSEVIER

Available online at www.sciencedirect.com

ScienceDirect

Current Opinion in
Microbiology

Unveiling new microbial eukaryotes in the surface ocean

Ramon Massana and Carlos Pedrós-Alió

A decade after molecular techniques were used to discover novel bacteria and archaea in the oceans, the same approach has revealed a wealth of new marine eukaryotic microbes. The approach has been particularly successful with the smallest eukaryotes, where morphological and culture approaches frequently fail. Analysis of samples from the surface ocean, the most accessible and supposedly well-known oceanic region, reveals novel eukaryotic diversity at all different levels: from the highest taxonomic rank to the lowest microdiverse clusters. Moreover, marine eukaryotic assemblages show a large diversity with members belonging to many different lineages. The implication of this large and novel eukaryotic diversity for biodiversity surveys and ecosystem functioning opens new avenues for future research.

Address

Department of Marine Biology and Oceanography, Institut de Ciències del Mar, CSIC, Passeig Marítim de la Barceloneta 37-49, 08003 Barcelona, Catalonia, Spain

Corresponding author: Massana, Ramon (ramonm@icm.csic.es)

Current Opinion in Microbiology 2008, **11**:213–218

This review comes from a themed issue on
Ecology and industrial microbiology
Edited by Juan Ramos and Martin Keller

Available online 13th June 2008

1369-5274/\$ – see front matter

© 2008 Elsevier Ltd. All rights reserved.

DOI [10.1016/j.mib.2008.04.004](https://doi.org/10.1016/j.mib.2008.04.004)

Introduction

Microorganisms are known to play essential roles in natural systems. In the surface oceans they are present in large abundances, account for a significant share of planktonic biomass, and are central in biogeochemical cycles. Yet, they are a heterogeneous collection of organisms with a substantial diversity of form, size, life style, and phylogenetic affiliation, and they are pooled together only because of their invisibility to the naked eye [1]. Many microorganisms, especially the smallest ones, cannot be identified under the microscope, because they lack conspicuous morphological features. This is well known in prokaryotes, which have only a handful of possible morphologies. Only after molecular surveys retrieving SSU rDNA sequences directly from the environment were carried out, the phylogenetic affiliation of marine prokaryotes was known. The first studies on the molecular diversity of marine bacteria [2] and archaea [3]

unveiled a substantial amount of new diversity and revealed that most microorganisms available in pure culture were not dominant in the marine plankton. Microbial eukaryotes, in turn, were not considered in these early approaches, despite the fact that the smaller eukaryotes were also difficult to be identified morphologically. In most conventional studies of marine plankton, these organisms were usually lumped in a black box labeled ‘small flagellates’. The growing recognition of the importance of these minute eukaryotes as primary producers, bacterial grazers, and parasites paralleled the interest of identifying the species of these probably very diverse assemblages. The first molecular surveys of marine eukaryotes confirmed the existence of significant novel diversity within the protistan world [4[•],5[•],6[•]]. This approach has benefited from recent efforts to build a robust framework of eukaryotic evolution within which environmental sequences can be placed. Virtually all eukaryotic organisms can be grouped into a few supergroups. Each one is composed by distinct lineages (most are protists) that are held together by phylogenetic signatures and some ultrastructural characters [7,8[•],9]. For instance, the supergroup opisthokonta includes metazoans, fungi, and choanoflagellates. Here we analyze and summarize the novel diversity of marine microbial eukaryotes as revealed by molecular studies in the surface of the oceans.

A new window into protistan diversity in the sea

Introduction of molecular tools in microbial ecology has become the key to access the phylogenetic and functional diversity of marine microbes [10]. The basis is to extract total DNA from a community, to amplify a marker gene (18S rDNA in eukaryotes) by PCR, and to clone and sequence the PCR products for phylogenetic identification. The first molecular surveys of marine eukaryotes targeted the smallest cells (picoeukaryotes, $\leq 3 \mu\text{m}$ in size) from surface [4[•],6[•]] and deep [5[•]] oceanic samples. Extreme environments, such as anoxic water column and sediments [11,12] and hydrothermal vents [13] were inspected soon after, in the search for the limits of eukaryotic life and the most divergent and ancient lineages. After the sequencing effort, specific oligonucleotide probes could be designed and applied through fluorescent *in situ* hybridization (FISH) to visualize the target cells in natural samples and to determine their distribution and abundance. This has been done for groups having both cultured [14] and uncultured [15,16[•],17] representatives. It is in the latter case where FISH exploits its full potential, allowing to put a face (cell size, rough shape, chlorophyll presence) to novel lineages

Table 1

Affiliation of 18S rDNA sequences from surface picoplankton in the coast (23 libraries and 1349 clones) and open sea (12 libraries and 826 clones) from studies that reported the clonal distribution among phylogenetic groups

Supergroup	First rank	Second rank	<i>n</i>	% clones	Habitat	Novelty ^a
Archaeplastida	Chloroplastida	Prasinophytae	261	15.0	Coast	Low
		Chlorophyta	2	0.1		–
Chromalveolata	Alveolates	Ciliophora	137	4.9	Coast	Medium
		Dinzoa	173	5.3		Low*
		MA-I	363	14.6		High
		MA-II	417	17.6		High
	Stramenopiles	Bicosoecida	20	0.4	Coast	Low
		Bolidophyceae	21	0.4		Low
		Chrysophyceae	65	2.7		Medium
		Diatoms	27	2.5		Low*
		Dictyochophyceae	25	1.1		Low
		Eustigmatales	2	0.4		–
		Labyrinthulids	14	0.5		Medium
		MASTs	237	13.4		High
		Oomycetes	8	0.2		–
		Pelagophyceae	12	1.0		Low
		Pirsonia	3	0.1		–
		Cryptophyceae	64	2.4		Low*
	Basal groups	Haptophyta	53	4.5	Offshore	Low*
		Katablepharis	6	0.2		–
		Telonema	14	0.4		Medium
Excavata	Euglenozoa	Kinetoplastea	1	0.0		–
Opisthokonta	Choanoflagellates		16	1.0		Medium
	Fungi		16	0.8		High
Rhizaria	Cercozoa		53	2.6	Coast	Medium
	Chlorarachniophyte-like		11	0.3		High
	Radiolaria		90	5.6		High
Not assigned	Apusomonads		11	0.2		Low
	Picobiliphyta		24	0.9		High
	Inserta sedis		29	0.7		High

Data derive from Pacific, Atlantic, Indian, and Southern Oceans, and the Mediterranean and North Seas [4^{••},6^{••},22–27]. The number of clones, the average contribution of each group (normalized by the number of clones in each system), the coast-offshore trend, and a rough indication of the level of novel diversity detected within the group are shown.

^a Low: most clones are ≥98% similar to cultured relatives (* indicates a significant presence of more divergent clones); medium: most clones are 92–98% similar to cultured relatives; high: most clones are <92% similar to cultured relatives.

detected by environmental sequencing. Finally, the metagenomic approach retrieves the full gene content of natural assemblages [18,19], providing a picture of microbial diversity independent both from culturing and PCR biases. This is done by direct cloning of small or large fragments of environmental DNA for later sequencing. Although metagenomic studies on marine microbes have so far targeted mostly the prokaryotic size fraction [20[•]], they contain a few 18S rDNA sequences [15]. A metagenomic study including picoeukaryotes has been recently published [21].

Here we analyze the 18S rDNA sequences of marine picoeukaryotes derived from the euphotic marine zone, the well-oxygenated surface skin of the ocean that is also most reactive to biogeochemical cycles. We examined 35 PCR-based libraries prepared with coastal and open sea samples from the Pacific [6^{••},22], Atlantic [23], Indian [24], and Southern Oceans [4^{••}], and the Mediterranean

[4^{••},25] and North Seas [26,27]. These are the libraries where enough information was provided to calculate relative clonal abundance, and yield information comparable to other studies [28–30]. The 2175 clones are distributed among all eukaryotic supergroups, with the exception of amoebzoa (Table 1). Alveolates (43% of clones), stramenopiles (23%), and prasinophytes (15%) are the better-represented groups. Some seem more important in coastal areas (cercozoans, ciliates, cryptophytes, alveolates-II, and prasinophytes), and others in the open sea (MASTs, pelagophytes, haptophytes, and radiolarians). Even in this supposedly well-known marine habitat, the most accessible and well studied, substantial novel diversity is revealed at different phylogenetic scales: high-rank novelty (sequences outside supergroups), intermediate-rank novelty (novel lineages within supergroups), and low-rank novelty (sequences close but not identical to characterized organisms, putatively representing new species, genera, orders, or families).

High-rank novel diversity: more than the six supergroups?

The first molecular surveys of marine protists claimed the discovery of novel groups that deserved the highest taxonomic rank, which could not be placed within any of the eukaryotic supergroups [5[•],11,12]. Many of these sequences derived from anoxic systems. However, it was soon shown that some of these highly divergent groups were unsupported because of the presence of undetected chimeras, misplacement of fast evolving lineages, and incomplete representation of cultured strains [31^{••},32,33]. However, several sequences still form robust and deep clades and thus remain as candidates for novel high-rank taxonomic groups. They are found at low clonal abundance so, probably, they are not very important ecologically. Instead, their interest is that they might represent different pathways in eukaryotic evolution. Perhaps the best example is the picobiliphytes, a novel phytoplanktonic class that is not close to any supergroup [17]. Picobiliphyte cells are 3–4 µm in size, putatively contain a plastid with phycobilin pigments, and can be locally abundant [15]. Other high-rank groups await careful scrutiny, because each library often yields a few sequences impossible to be classified to a given supergroup. Twenty-nine clones of the overview presented here could not be assigned to a given supergroup (Table 1). The similarity of these sequences is often below 75% to any other known 18S rDNA. Some of them appear in more than one system, thus excluding the possibility of being chimeras. Exhaustive phylogenetic reconstructions, renovated culturing efforts, and additional sequences from single cells are needed to understand the biological nature of these putative high-rank lineages.

Intermediate-rank novel diversity: novel eukaryotic lineages

A large number of sequences form clades that affiliate to a given eukaryotic supergroup but without a clear affiliation to any defined group. Among these, the marine alveolates (MAs) and marine stramenopiles (MASTs) are particularly interesting because they appear in virtually all marine surveys. MAs are divided into two main groups, MA-I and MA-II, which form robust lineages equivalent to the other alveolate groups. Their placement is still unresolved, because SSU analysis places them closer to dinoflagellates [34], whereas a recent LSU tree places them closer to perkinsids [21]. The sequence diversity contained in the two groups is huge, with at least 5 clades within MA-I and up to 16 clades within MA-II [35[•]]. Besides their genetic diversity, these groups appear at a very high clonal abundance, 14.6% for MA-I and 17.6% for MA-II. Soon after the description of MAs, *Amoebophrya* (a dinoflagellate parasite) was sequenced and affiliated to MA-II [36]. Additional sequences of parasites have been later published within both MA-I and MA-II [34,37,38]. Therefore, it has been proposed that the whole

assemblage is composed of parasites of marine organisms. Perhaps, the specific interaction with different hosts could explain their large genetic diversity. Their considerable clonal abundance and diversity suggests an important role for parasitism as a trophic relationship in the open sea.

MASTs form more than 10 clades at the basal part of the stramenopiles [39[•]], where all protists are heterotrophic, including free-living phagotrophic flagellates (bicosoecids), parasites (*Blastocystis*), or osmotrophs (oomycetes and labyrinthulids). MASTs are rather abundant (13.4% of clones), and a few clades (MAST-1, MAST-3, MAST-4, and MAST-7) account for most sequences (the other clades have lower clonal abundances or are specific of anoxic systems). The heterotrophic nature of MASTs, first suspected by their phylogenetic placement, was confirmed by FISH for clade-1, clade-2, and clade-4 [16^{••}]. These are small protists (2–8 µm in size), able to grow in the dark and to ingest bacteria. These MAST cells are widely distributed and account for a significant fraction of heterotrophic flagellates globally. One group in particular, MAST-4, is found in all samples (except the polar ones) as a very small protist (2–3 µm in size), its abundance averages 130 cells ml⁻¹, and accounts for 9% of heterotrophic flagellates. Overall, these results reveal that still-uncultured groups can be dominant in the oceans and highlight the ecological relevance of the novel diversity detected by the molecular approach.

Low-rank novel diversity: known lineages are more diverse than thought

This is represented by sequences that clearly affiliate to a given lineage but are not identical to any characterized protist. Low-rank novel diversity is extensive, because environmental sequences are identical to cultured strains only in a few cases. For instance, only 11 out of 510 partial sequences retrieved from the Indian Ocean [24] were identical (over 800–900 bp) to cultured strains: *Caecitellus parvulus*, *Micromonas pusilla*, and *Ostreococcus* RCC 143. This list would include *Bathycoccus prasinos*, *Amastigomonas debruynei*, *Gymnodinium* sp., and *Pelagomonas calceolata* if 1–2 mismatches were accepted (another 21 clones). So, between 94 and 98% of the sequences retrieved from this marine system represent new diversity not explained by cultured protists.

Obviously, there are degrees of novelty at this lower phylogenetic level. Thus, prasinophytes show the best correspondence between molecular and culturing approaches, and the 18S rDNA sequences from the field and cultures are identical or very close [40]. Other important groups often represented by environmental sequences closely related to cultured strains are the bicosoecids, bolidophytes, dinoflagellates, and pelagophytes. These closely related sequences most probably

identify the same cultured strain in the environment, closely related species, or ecotypes of the same species.

Other groups contain a much larger level of diversity. So, important marine groups such as the ciliates, choanoflagellates, chrysophytes, cryptophytes, diatoms, and haptophytes are represented by environmental sequences that can range from 94 to 100% sequence similarity to cultured relatives. This implies that some novel clades exist within all these groups, increasing substantially the diversity they contain. For instance, up to three novel clades have been identified within the choanoflagellates, and three novel clades within chrysophytes after analyzing marine sequences (del Campo, Massana, unpublished).

A substantial fraction of clones from the open sea affiliate with the radiolarians (10% on average). This is surprising, because the radiolarian species known so far are rather large (typically around 100 μm) and most possess mineralized skeletons. Marine radiolarian sequences are diverse, generally highly distant to sequenced protists, and form at least five clades, two related to acanthareans, one to polycystinea, and two to taxopodida [23]. Whereas it is known that many described radiolarian species have not been sequenced, given the difficulty of their isolation, it is not clear if these would explain the sequences found in environmental surveys. The existence of these diverse radiolarian sequences from the picoplankton remains as an intriguing enigma.

Microdiversity of natural assemblages

Prokaryotic diversity is normally structured in clades containing highly related but seldom identical sequences [41,42]. The evolutionary and ecological meaning of this microdiversity is not well understood, though it has been proposed that it is a consequence of the asexual mode of prokaryotic reproduction together with ecological factors [43]. Microdiverse clusters would exist because of neutral mutations (also in the 18S rDNA) during asexual divisions, so that all the members would occupy the same ecological niche. When one of these members acquires a selective advantage, periodic selective sweeps would purge all variability within the cluster. Thus, current microdiverse clusters would exist due to the accumulation of neutral mutations since the last selective sweep [44]. It is not clear whether this scenario also holds for microbial eukaryotes. First, it has to be demonstrated that microbial eukaryotes show microdiversity in nature. Some data suggest that this may be the case, at least for some groups such as MASTs [39^{*}] and MAs [35^{*}]. Second, even though cell division in microbial eukaryotes is mostly asexual, it is known (at least for some groups) that sexual events also occur, and these would certainly impact the genetic structure of populations by making populations more homogeneous. It is improbable that sexually compatible organisms show any variation within the 18S rDNA.

Mating experiments with related diatom strains showing some genetic structure reveal that only those with an identical 18S rDNA are sexually compatible [45^{**}]. If this applies to other marine populations, then each different 18S rDNA sequence, even with a single base pair difference, would mean an independent and evolutionarily isolated lineage, increasing protist diversity enormously. Thus, the actual microdiversity structure within particular protistan lineages deserves a better study, together with studies to unravel the sexual nature of marine protists.

Conclusions

The analysis of the 18S rDNA sequences retrieved from the sea reveals that marine protists are very diverse, increasing substantially the known amount of diversity within the eukaryotic tree of life. Current parametric and nonparametric estimates of protistan richness [46] indicate that hundreds to thousands of distinct protistan taxa can coexist in a single marine sample [25,28,29]. Most probably, microorganisms do not deviate from the trend of increasing number of species with decreasing individual size [47,48]. As shown here, the relatively low number of microbial species actually described is largely because of the under-representation of microbial diversity in culture collections. The availability of powerful and relatively cheap sequencing techniques will be essential to determine the dimensions of such diversity.

Another interesting point is that the increase in marine eukaryotic diversity occurs at almost all possible phylogenetic scales. Thus, putative high-rank groups occur, novel clades within supergroups have been identified, and novel diversity is detected within all known lineages, from closely new species, genera, families, or orders. The challenge is to retrieve in culture the organisms responsible for such sequences and to determine their trophic role and ecological function.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Sherr EB, Sherr BF: **Marine microbes. An overview.** In *Microbial Ecology of the Oceans*. Edited by Kirchman DL. New York: Wiley-Liss; 2000:13-46.
This book chapter is a good introduction to the main groups of marine microorganisms, organized by sizes, classes, general taxonomic assemblages, and functional groups.
2. Giovannoni SJ, Britschgi TB, Moyer CL, Field KG: **Genetic diversity in Sargasso Sea bacterioplankton.** *Nature* 1990, **345**:60-63.
3. DeLong EF: **Archaea in coastal marine environments.** *Proc Natl Acad Sci U S A* 1992, **89**:5685-5689.
4. Díez B, Pedrós-Alió C, Massana R: **Study of genetic diversity of eukaryotic picoplankton in different oceanic regions by small-subunit rRNA gene cloning and sequencing.** *Appl Environ Microbiol* 2001, **67**:2932-2941.

This is one of the first studies that report marine picoeukaryotic diversity by analyzing 18S rDNA genes directly retrieved from the environment. It compares data from distant surface waters and derives the general patterns of protistan diversity: high diversity and novel groups.

5. López-García P, Rodríguez-Valera F, Pedrós-Alió C, Moreira D:

- **Unexpected diversity of small eukaryotes in deep-sea Antarctic plankton.** *Nature* 2001, **409**:603-607.

This is one of the first studies that report marine picoeukaryotic diversity by sequencing environmental 18S rDNA genes, with special focus on deep oceanic samples as examples of extreme and undersampled systems.

- 6. Moon-van der Staay SY, De Wachter R, Vaulot D: **Oceanic 18S rDNA sequences from picoplankton reveal unsuspected eukaryotic diversity.** *Nature* 2001, **409**:607-610.

This is one of the first studies that report marine picoeukaryotic diversity by sequencing environmental 18S rDNA genes from a very oligotrophic oceanic sample.

- 7. Adl SM, Simpson AGB, Farmer MA, Andersen RA, Anderson OR, Barta JR, Bowser SS, Brugerolle G, Fensome RA, Fredericq S *et al.*: **The new higher level classification of eukaryotes with emphasis on the taxonomy of protists.** *J Eukaryot Microbiol* 2005, **52**:399-451.

- 8. Baldauf SL: **The deep roots of eukaryotes.** *Science* 2003, **300**:1703-1706.

This article is one of the first efforts to summarize all eukaryotic diversity into a few supergroups (eight in this study) using the information gathered in multigene phylogenetic trees.

- 9. Simpson AGB, Roger AJ: **The real "kingdoms" of eukaryotes.** *Curr Biol* 2004, **14**:R693-R696.

- 10. Pace NR: **A molecular view of microbial diversity and the biosphere.** *Science* 1997, **276**:734-740.

- 11. Dawson SC, Pace NR: **Novel kingdom-level eukaryotic diversity in anoxic environments.** *Proc Natl Acad Sci U S A* 2002, **99**:8324-8329.

- 12. Stoeck T, Epstein S: **Novel eukaryotic lineages inferred from small-subunit rRNA analyses of oxygen-depleted marine environments.** *Appl Environ Microbiol* 2003, **69**:2657-2663.

- 13. Edgcomb VP, Kysela DT, Teske A, de Vera Gomez A, Sogin ML: **Benthic eukaryotic diversity in the Guaymas Basin hydrothermal vent environment.** *Proc Natl Acad Sci U S A* 2002, **99**:7658-7662.

- 14. Not F, Latasa M, Marie D, Cariou T, Vaulot D, Simon N: **A single species, *Micromonas pusilla* (Prasinophyceae), dominates the eukaryotic picoplankton in the Western English Channel.** *Appl Environ Microbiol* 2004, **70**:4064-4072.

- 15. Cuvelier M, Ortiz A, Kim E, Moehlig H, Richardson DE, Heidelberg JF, Archibald JM, Worden AZ: **Widespread distribution of a unique marine protistan lineage.** *Environ Microbiol* 2008, **10**:1621-1634.

- 16. Massana R, Terrado R, Forn I, Lovejoy C, Pedrós-Alió C: **Distribution and abundance of uncultured heterotrophic flagellates in the world oceans.** *Environ Microbiol* 2006, **8**:1515-1522.

This article applies FISH to visualize uncultured protists and report their cell size and abundance. These are globally distributed heterotrophic flagellates with a significant abundance.

- 17. Not F, Valentin K, Romari K, Lovejoy C, Massana R, Töbe K, Vaulot D, Medlin L: **Picobiliphytes: a marine picoplanktonic algal group with unknown affinities to other eukaryotes.** *Science* 2007, **315**:252-254.

- 18. Venter JC, Remington K, Heidelberg JF, Halpern AL, Rusch D, Eisen JA, Wu D, Paulsen I, Nelson KE, Nelson W *et al.*: **Environmental genome shotgun sequencing of the Sargasso Sea.** *Science* 2004, **304**:66-74.

- 19. DeLong EF, Preston CM, Mincer T, Rich V, Hallam SJ, Frigaard N-U, Martinez A, Sullivan MB, Edwards R, Rodriguez Brito B *et al.*: **Community genomics among stratified microbial assemblages in the ocean's interior.** *Science* 2006, **311**:496-503.

- 20. Rusch DB, Halpern AL, Sutton G, Heidelberg KB, Williamson S, Yooseph S, Wu D, Eisen JA, Hoffman JM, Remington K *et al.*: **The Sorcerer II global ocean sampling expedition: northwest**

Atlantic through eastern tropical Pacific. *PLoS Biol* 2007, **5**:398-431.

This article presents a paramount amount of genetic information from surface marine microorganisms by doing a systematic metagenomic survey across a wide oceanographic transect.

- 21. Massana R, Karniol B, Pommier T, Bodaker I, Béjà O: **Metagenomic retrieval of a ribosomal DNA repeat array from an uncultured marine alveolate.** *Environ Microbiol* 2008, **10**:1335-1343.

- 22. Worden AZ: **Picoeukaryote diversity in coastal waters of the Pacific Ocean.** *Aquat Microb Ecol* 2006, **43**:165-175.

- 23. Not F, Gausling R, Azam F, Heidelberg JF, Worden AZ: **Vertical distribution of picoeukaryotic diversity in the Sargasso Sea.** *Environ Microbiol* 2007, **9**:1233-1252.

- 24. Not F, Latasa M, Scharek R, Viprey M, Karleskind P, Balagué V, Ontoria I, Cumino A, Goetze E, Vaulot D *et al.*: **Protistan assemblages across the Indian Ocean, with a specific emphasis on the picoeukaryotes.** *Deep Sea Res*, in press.

- 25. Massana R, Balagué V, Guillou L, Pedrós-Alió C: **Picoeukaryotic diversity in an oligotrophic coastal site studied by molecular and culturing approaches.** *FEMS Microbiol Ecol* 2004, **50**:231-243.

- 26. Medlin LK, Metfies K, Mehl H, Wiltshire K, Valentin K: **Picoeukaryotic plankton diversity at the Helgoland time series site as assessed by three molecular methods.** *Microb Ecol* 2006, **52**:53-71.

- 27. Romari K, Vaulot D: **Composition and temporal variability of picoeukaryote communities at a coastal site of the English Channel from 18S rDNA sequences.** *Limnol Oceanogr* 2004, **49**:784-798.

- 28. Countway PD, Gast RJ, Savai P, Caron DA: **Protistan diversity estimates based on 18S rDNA from seawater incubations in the western North Atlantic.** *J Eukaryot Microbiol* 2005, **52**:95-106.

- 29. Countway PD, Gast RJ, Denmet MR, Savai P, Rose JM, Caron DA: **Distinct protistan assemblages characterize the euphotic zone and deep sea (2500 m) of the western North Atlantic (Sargasso Sea and Gulf Stream).** *Environ Microbiol* 2007, **9**:1219-1232.

- 30. Lovejoy C, Massana R, Pedrós-Alió C: **Diversity and distribution of marine microbial eukaryotes in the Arctic Ocean and adjacent seas.** *Appl Environ Microbiol* 2006, **72**:3085-3095.

- 31. Berney C, Fahrni J, Pawlowski J: **How many novel eukaryotic "kingdoms"? Pitfalls and limitations of environmental DNA surveys.** *BMC Biol* 2004, **2**:1-13.

This article performs a rigorous check on the claims of novel high-rank diversity in the initial molecular surveys. Some sequences were shown to be artifacts, but candidates for high-rank diversity still remain.

- 32. Cavalier-Smith T: **Only six kingdoms of life.** *Proc R Soc Lond B* 2004, **271**:1251-1262.

- 33. Epstein S, López-García P: **"Missing" protists: a molecular prospective.** *Biodivers Conserv* 2008, **17**:261-276.

- 34. Skovgaard A, Massana R, Balagué V, Saiz E: **Phylogenetic position of the copepod-infesting parasite *Syndinium turbo* (Dinoflagellata, Syndinea).** *Protist* 2005, **156**:413-423.

- 35. Groisillier A, Massana R, Valentin K, Vaulot D, Guillou L: **Genetic diversity and habitats of two enigmatic marine alveolate lineages.** *Aquat Microb Ecol* 2006, **42**:277-291.

This article summarizes and labels the genetic diversity within uncultured marine alveolates (MAs) retrieved from marine picoeukaryotes.

- 36. Gunderson JH, John SA, Boman WC II, Coats DW: **Multiple strains of the parasitic dinoflagellate *Amoebophrya* exist in Chesapeake Bay.** *J Eukaryot Microbiol* 2002, **49**:469-474.

- 37. Dolven JK, Lindqvist C, Albert VA, Björklund KR, Yuasa T, Takahashi O, Mayama S: **Molecular diversity of alveolates associated with neritic North Atlantic radiolarians.** *Protist* 2007, **158**:65-76.

- 38. Harada A, Ohtsuka S, Horiguchi T: **Species of the parasitic genus *Duboscquella* are members of the enigmatic marine alveolate group I.** *Protist* 2007, **158**:337-347.

39. Massana R, Castresana J, Balagué V, Guillou L, Romari K, Groisillier A, Valentin K, Pedrós-Alió C: **Phylogenetic and ecological analysis of novel marine stramenopiles**. *Appl Environ Microbiol* 2004, **70**:3528-3534.

This article summarizes and labels the genetic diversity within uncultured marine stramenopiles (MASTs) retrieved from marine picoeukaryotes.

40. Guillou L, Eikrem W, Chrétiennot-Dinet M-J, Le Gall F, Massana R, Romari K, Pedrós-Alió C, Vaulot D: **Diversity of picoplanktonic Prasinophyceae assessed by direct SSU rDNA sequencing of environmental samples and novel isolates retrieved from oceanic and coastal marine ecosystems**. *Protist* 2004, **155**:193-214.
41. Acinas SG, Klepac-Ceraj V, Hunt DE, Pharino C, Ceraj I, Distel DL, Polz MF: **Fine-scale phylogenetic architecture of a complex bacterial community**. *Nature* 2004, **430**:551-554.
42. Fuhrman JA, Campbell L: **Microbial microdiversity**. *Nature* 1998, **393**:410-411.
43. Cohan FM: **Towards a conceptual and operational union of bacterial systematics, ecology, and evolution**. *Philos Trans R Soc B* 2006, **361**:1985-1996.

44. Giovannoni SJ: **Oceans of bacteria**. *Nature* 2004, **430**:515-516.

45. Amato A, Kooistra WHCF, Ghiron JHL, Mann DG, Pröschold T, Montresor M: **Reproductive isolation among sympatric cryptic species in marine diatoms**. *Protist* 2007, **158**:193-207.

This article compares the genetic distance, ultrastructure, and mating capacity of several diatom strains from the same species, showing a remarkable consistency among the three identifying properties.

46. Jeon S-O, Bunge J, Stoeck T, Barger KJ-A, Hong S-H, Epstein SS: **Synthetic statistical approach reveals a high degree of richness of microbial eukaryotes in an anoxic water column**. *Appl Environ Microbiol* 2006, **72**:6578-6583.
47. Hughes Martiny JB, Bohannan BJM, Brown JH, Colwell RK, Fuhrman JA, Green JL, Horner-Devine MC, Kane M, Krumins JA, Kuske CR *et al.*: **Microbial biogeography: putting microorganisms on the map**. *Nat Rev* 2006, **4**:102-112.
48. May RM: **How many species are there on Earth?** *Science* 1988, **241**:1441-1449.